



OMPHALINA

ISSN 1925-858

FORAY 2015 REPORT

Newsletter of



Vol. VI , No 7
Dec. 10, 2015



FORAY NEWFOUNDLAND AND LABRADOR

is an amateur, volunteer-run, community, not-for-profit organization with a mission to organize enjoyable and informative amateur mushroom forays in Newfoundland and Labrador and disseminate the knowledge gained.

Webpage: www.nlmushrooms.ca

ADDRESS

Foray Newfoundland & Labrador
21 Pond Rd.
Rocky Harbour NL
A0K 4N0
CANADA

E-mail: [info AT nlmushrooms DOT ca](mailto:info@nlmushrooms.ca)

BOARD OF DIRECTORS

Michael Burzynski

PRESIDENT

Geoff Thurlow

TREASURER

Tina Leonard

SECRETARY

André Arsenault

Jim Cornish

Jamie Graham

Tina Leonard

Anne Marceau

Erin McKee

Robert MacIsaac

Faye Murrin

Michele

Piercey-Normore

Helen Spencer

Marian Wissink

CONSULTANTS

MYCOLOGICAL

Dave Malloch

NB MUSEUM

AUDITOR

Gordon Janes

BONNELL COLE JANES

LEGAL COUNSEL

Andrew May

BROTHERS & BURDEN

OMPHALINA is the lackadaisical newsletter of Foray Newfoundland & Labrador. There is no schedule of publications, no promise to appear again. Its primary purpose is to serve as a conduit of information to registrants of the upcoming foray and secondarily as a communications tool with members.

Issues of **OMPHALINA** are archived in:

Library and Archives Canada's Electronic Collection <<http://epe.lac-bac.gc.ca/100/201/300/omphalina/index.html>>, and

Centre for Newfoundland Studies, Queen Elizabeth II Library, where a copy is also printed and archived <<http://collections.mun.ca/cdm4/description.php?phpReturn=typeListing.php&id=162>>.

The content is neither discussed nor approved by the Board of Directors. Therefore, opinions expressed do not represent the views of the Board, the Corporation, the partners, the sponsors, or the members. Opinions are solely those of the authors and uncredited opinions solely those of the Editor.

Please address comments, complaints and contributions to the largely self-appointed Editor, Andrus Voitk:

seened AT gmail DOT com,

*... who eagerly invites contributions to **OMPHALINA**, dealing with any aspect even remotely related to mushrooms. Authors are guaranteed instant fame—fortune to follow. Authors retain copyright to published material, and submission indicates permission to publish, subject to the usual editorial decisions. Issues are freely available to the public on the FNL website. Because content is protected by authors' copyright, editors of other publications wishing to use any material, should ask first.*

COVER: The cliffs of Western Brook Pond,

Gros Morne National Park.

Photo by: Michael Burzynski

CONTENTS

| | |
|---|-------------------|
| Message from the editor | 2 |
| Words from the president | 3 |
| Faculty and participants | 4 |
| Life at the edge; An arctic-alpine collecting trip on Big Level. | |
| <i>Michael Burzyki</i> | 6 |
| Four lichenomphalias from the top of Big Level | |
| <i>Andrus Voitk</i> | 8 |
| Tablelands lichen survey update | |
| <i>Michele Piercey-Normore</i> | 10 |
| <i>Russula griseascens</i> ; project update | |
| <i>Maria Voitk</i> | 13 |
| Foray fotos | 14 |
| Program | 20 |
| Workshops and walks | 22 |
| Trails | 29 |
| SPECIES LIST AND DISTRIBUTION BY FORAY TRAIL | |
| <i>Andrus Voitk, Michael Burzynski, Tony Wright</i> | 30 |
| What do these data mean? | |
| <i>Andrus Voitk</i> | 42 |
| SPECIES LIST AND DISTRIBUTION BY FORAY TRAIL - LICHENS | |
| <i>Chris Deduke, Michele Piercey-Normore</i> | 46 |
| Survey of the lichen-forming ascomycetes during the 2015 NL Foray | |
| <i>Chris Deduke, Michele Piercey-Normore</i> | 50 |
| Group photo 2015 | 54 |
| Our partner organizations | inside back cover |
| Notice - Goose Bay 2016 | back cover |

Message from the Editor

It is that time of year again to get the foray report out the door and onto your screens before too much snow flies and planning starts in earnest for the next one. This report details another successful foray in beautiful Gros Morne National Park. We were lucky, once again, to have a fabulous group of participants and experts come together in search and study of the region's fungi and lichens.

This report exists because of the hard work of many folks. They helped by writing articles, sending photos, proofreading, and by compiling and studying the results. Without their dedicated efforts, there would be no report or likely, no foray.

To all of you, participants, trail leaders, data base folks, faculty, organisers, a big thank you!

Wishing you all the best of the holiday season and looking forward to seeing you at next year's foray.

Marian Wissink
Guest Editor



Words from the President

Numbers always make me drowsy, and as usual the foray report is full of numbers. But the numbers are one of the big reasons that we do this fungus and lichen survey every year—they condense the information about the foray into a concise form, and let us compare forays from year to year and from place to place.

This year we had 58 participants, including nine faculty. This means that there were a lot of both experienced and fresh eyes looking for specimens—and these are the most important tools that we use at each foray. Without your help, very few of these species would have been found. Once again, Killdevil Camp and Gros Morne National Park exceeded our expectations. Foray 2015 was our fifth foray in Gros Morne. The area, with the national park and Sir Richard Squires Provincial Park, contains a wide diversity of habitats easily accessible by trails. Of the thousands of fungi and lichens collected, 1,575 were identified and databased—representing 397 taxa, Roger took 710 specimen photographs (his trigger finger is still smoking), and 986 dried specimens were added to our fungarium. As Andrus points out in his analysis of the list, we are still far from exhausting the potential of this province's mycota.

During the pre-foray field trips with the faculty, Parks Canada gave us the opportunity to helicopter eight people onto the Long Range to sample the tundra overlooking Western Brook Pond. There, at the top of the 600 metre cliffs, we spent four hours collecting fungi and lichens—the first time that we have been able to survey the park highlands. Interestingly, in terms of species the highlands were the third most productive habitat surveyed during the foray.

I would like to express my thanks to this year's experts for offering their expertise and time; to Mac Turner and Killdevil Camp for providing us with wonderful accommodations, work spaces, and meals; to Superintendent Geoff Hancock and Parks Canada for erecting the sorting tent and offering helicopter support; to Chef Chris Chaisson and his crew for the fantastic Saturday cookup; to Chef Jason Nesbitt for the wonderful fungus-themed Faculty Dinner on Sunday evening, and to everyone who volunteered their effort and eyeballs to make this foray work.

Now I'll just close *my* eyes for a while until the numbers stop swirling around.

Michael





FACULTY

Guest faculty:

Oluna Ceska
Nils Hallenberg
Andy Methven
Andrew Miller
Michele Piercey-Normore
Greg Thorn
Roger Smith

Local Faculty:

Michael Burzynski
Faye Murrin
Andrus Voitk



PARTICIPANTS

Chris Deduke
Bruce Pendergast
Barbara Pendergast
Mildred Brown
Tony Dearness
Michel Savard
Isabella St. John
Glynn Bishop
Tarik Rodrigues
Bruce Rodrigues
Kaden Rodrigues
Andrus Voitk
Maria Voitk
Serena Hancock
David Bazeley
John Burghardt
Jenifer Burghardt
Peter Thurlow
Meigan Aspin
Nicholas Michalski
Michele Piercey-Normore
Ryan Haley
Robert MacIsaac
Carolyn Molson
Michael Burzynski
Anne Marceau
Anthony E. McNally
Nils Hallenberg
Roger Smith
Verlé Harrop

Christopher Chaisson
Jamie Graham
Tina Leonard
Tony Wright
Faye Murrin
Helen Spencer
Don Spencer
Roger Zilkowsky
Graham Zilkowsky
André Arsenault
Marian Wissink
Jason Nesbitt
Claudia Hanel
Shane White
Elaine Humber
Yolanda Weirsmas
John Sandlos
William
Xavier
Hilary Margaret Harrop-Archibald
Milena Orchard-Hays
Greg Thorn
Adolf Ceska
Oluna Ceska
Andy Methven
Andy Miller
Jessica Butler
Pieter van Heerden
Katherine Flores

Photo: TA Loeffler

Life at the Edge

An Arctic-alpine Collecting Trip on Big Level

Michael Burzynski



Each foray is preceded by several days of faculty foray when visiting mycologists and lichenologists become familiar with the area and collect species of interest to them. We try to take an ecosystem approach with our sampling, and this year we had the opportunity to explore a habitat that we had not previously been able to reach—the highlands of the Long Range Mountains.

The highest point in Gros Morne is 806 metres above sea level, but strong winds and coastal climate force tree level down to only 600 metres. Above that is a landscape of arctic-alpine heath, rock barrens, and impenetrable stunted spruce-fir forest known locally as tuckamore.

During Faculty Foray 2015, Parks Canada offered us the chance to helicopter eight participants onto the highlands of Gros Morne National Park and to spend four hours sampling the Arctic-alpine Zone in the Big Level area, just south of Western Brook Pond. This is a wilderness area almost at the centre of the park, and there are no trails or other means of access. The team consisted of Greg Thorn, Michele Piercey-Normore, Chris Deduke, Faye Murrin, Andrus and Maria

Voitk, Anne Marceau, and Michael Burzynski. Andrus and Maria concentrated on the tiny omphalinoids (see Andrus' article in this issue), Michele and Chris sampled lichens throughout the area, and the rest of us collected whatever we could find.

On the highlands, size matters. If you are a plant and you get too tall, cold dry winter winds will cut you back down to size. Because of this, hollows and gullies are a tangle of dwarfed trees, and exposed plateaus are usually lichen-covered outcrops surrounded by ankle-high shrubs. Mushrooms are some of the tallest features of the heathland, since they do not have to survive through the winter—the actual fungus organisms are protected within the soil. Many of the largest mushrooms in the heath and tuckamore are the typical species associated with trees and shrubs on the lowlands. On the highlands, these mushrooms seem outsized because they dwarf the surrounding vegetation.

By crawling into the tuckamore tangle we collected wood-decaying corticioids for Nils to work with back at the lab. He found that many of the species on the highlands are very different from those in the

nearby lowland forest. This is an area that we should explore more thoroughly in the future.

This was a very productive trip, and a wonderful chance to start an inventory of the fungi of the highlands. Please see the Big Level column of the species list in this issue of *Omphalina* for a full tally of what was collected, and our *Flickr* site for photographs. We are indebted to Field Unit Superintendent Geoffrey Hancock and Ecosystem Scientist Tom Knight for making our Long Range sampling possible.

Anne spotted this Black Bear watching us from the edge of the cliff.





Some fungi were taller than the surrounding vegetation, but the tiny omphalinoids required a close search.



Andrus and the helicopter pilot. Our flights gave us great views of moose-altered forest, the boggy coastal lowlands, and the uplifted Long Range plateau. Photo Greg Thorn.



Michele sampling crustose lichens. Photo Greg Thorn.



Greg searching the tuckamore forest.



Michael photographing the gorge of Western Brook Pond. The area that we sampled was about 640 metres above the water. Photo Greg Thorn.

Four lichenomphalias from the top of Big Level

Andrus Voitk



Photo: Maria Voitk

There is no question that for me the trip to Big Level during the Faculty Foray was the highlight of this year's foray. The sparse and barren arctic-alpine habitat speaks directly to my heart like no other. Partial to this habitat, I am drawn to the animals, plants and fungi that live there; my favourites are the small lichenomphalias for whom this is home. The title banner photo shows prime real estate for these species, and sure enough, we found all four species known to exist in this province within steps of this site.

In 2006 I wrote about the three *Lichenomphalia* species we had seen in the province, after finding all three on top of Gros Morne Mountain.¹ In 2011 Maria and I discovered a small brown mushroom atop Mt Ignoble near our home,² but it took us another few years to recognize that there were more than one species there, and only one of these was a

Lichenomphalia. Listed as an *Omphalina* for now, its exact identity is still being investigated: in the previous issue we described our attempt to locate its topotype on Mt Washington.³ Well, of the four *Lichenomphalia* species on Big Level at the time, this small brown mushroom was the commonest. These finds will contribute greatly to the eventual accurate identification of this species.

Most lichens are ascomycetes. These four are lichenized (live mutualistically with algae) agaric (cap, gills and stem) basidiomycetes. Algal granules can be seen at the base of the stem for three; one, *L. hudsoniana*, has a unique leafy lichen thallus, that identifies it even in the absence of the fruiting body.

These four species are illustrated on the next page with photographs from Big Level. Beautiful subjects make even mediocre photos look gorgeous.

References

1. Voitk A, Voitk M, Mann H, Lebeuf R: *Arhenia obscurata*—the phoenix of Mt Ignoble. *OMPHALINA* 5(7):5-21. 2014.
2. Voitk A: Three lichenomphalias from the top of Gros Morne Mountain. *The Mycophile* 47(5):1&11. 2006.
3. Voitk A, Voitk M: The elusive fairy. *OMPHALINA* 6(6):15-19. 2015.

Illustrations: **A** *Lichenomphalia alpina*. Yellow cap and yellow stem, green mat of algal granules at base. **B** *Lichenomphalia hudsoniana*. Yellow cap, white stem, algae in leafy lichen thallus at base. **C** *Omphalina* cf. *oreades*. Brown small mushroom with hairy stem, green mat of algal granules at base. **D** *Lichenomphalia umbellifera*. White to tan mushroom, mat of algal granules at base. Photo not the best, so trust me!

Title banner: Arctic-alpine heath on the edge of the Western Brook Pond gorge. The Pond is over 500 m straight down—the tour boat looked like a very small toy making its way silently across the surface.



Photo: Maria Voith

Tablelands lichen survey update

Michele Piercey-Normore

The serpentine rock of the Tablelands was heaved to the surface from the Earth's mantle during collision of tectonic plates about 500 million years ago. Full of chromium, iron and nickel, they are inhospitable to the growth of most organisms (background photo). Therefore, the few lichens that do grow on them (upper insert) are of special interest, because of their unusual preference, adaptation or tolerance.

Among the serpentine rocks are also erratics (middle insert), so called because they were dropped here by movement of water and ice during times of glaciation. Their more user friendly composition permits ready colonization by lichens, making their colourful presence obvious among the barren serpentine moonscape.

Particles from crumbling rocks accumulate in cracks between them, which also trap some wind-blown organic debris over the years, providing a substrate for hardier organisms. Lichens grow on this soil and its plants (lower insert). These areas are particularly rich in border regions of advancing heath and woodland.

Our lichen survey of the Tablelands, although particularly interested in documenting the species that grow on serpentine, also reviews lichens from the other two zones.

Photos: Michael Burzynski & Roger Smith



The lichen flora on the serpentine Tablelands in Gros Morne National Park was investigated during two days last Fall (September 19-20, 2015) with the permission of the park. While lichen surveys have been conducted on peridotite rock of other major world sites, this will be the first lichen survey for the serpentine fields in Newfoundland despite this area having been declared a World Heritage Site. The goal of our project was to collect representatives of all species in the area. Four locations along the south side of Highway 431 were chosen beginning with the Tablelands trail to explore the serpentine barrens, which were predominantly peridotite rock and a few erratics deposited from glaciation. Three additional sites between the trail and Trout River were all on serpentine but they contained an increasing number of erratics further west of the Tablelands Trail

because of glacial debris carried from elsewhere. These sites were colloquially called Halfway Cliff, the wind tunnel, and the serpentine opposite Wallace Brook. By sampling from all types of rocks, soil, vegetation, and other debris we hope to determine whether the species on peridotite are unique to the peridotite rock or whether they are present on all types of rocks and can tolerate the conditions of the peridotite rocks.

The participants in the trip included Michael Burzynski, Anne Marceau, who were collectors and guides for the area; Joe Brazil, who was our designated photographer; and Chris Deduke and myself who collected lichens. We were joined by two people from Parks Canada [Kris Oravec and Erin McKee] for a short time on the first day who




Photo: Joe Brazil

Michael Burzynski on a large boulder of peridotite. In the foreground peridotite rubble and very little vegetation. This is contrasted with a background of forest.



This one we called the Ghost lichen because the centre of the lichen was always absent and the only part visible was the younger ring. This was one of the two lichens consistently found on peridotite.

explored the flora and geology in the area; and Roger Smith for the second day who photographed the diversity and landscape. Over the course of two days more than 200 specimens were collected, which were obtained by chipping small pieces of rock with a hammer and chisel, scraping lichens from rock, bark, and soil, and sometimes having to carry larger rocks back for lichen identification. So far two species have been determined to be common on the peridotite rock while several other less frequent species are also present. Other lichens on different types of substrates were diverse and include over 70 species waiting for further confirmation. Explorations of the three

other major serpentine mountains in the province are planned for the next three years including the Lewis Hills, the Blowmedowns, and the Gregory Plateau. By the end of four years we plan to have a better understanding of the lichens that can grow in these inhospitable conditions and will be able to compare our serpentine lichen flora with that of other serpentine areas in the world.



RUSSULA GRISEASCENS

project update

Maria Voltk

You may remember that *OMPHALINA* 6(1) dealt with a global fungal survey using soil samples. One unexpected result of our contributions was to find no DNA of the common *Russula emetica*

in our soil, but abundant samples close to *R. griseascens*, a species we had never identified before. After a review and some consultations and discussions, we concluded that perhaps until things get sorted out, we might be more correct to apply the name *R. griseascens* to what we had been calling *R. emetica* to date.

Well, up stepped Anna Bazzicalupo from BC, who has looked at the *R. emetica* complex in the Pacific Northwest, with an offer to study it here for a continental picture. She sent along vials of buffer to keep the DNA safe, and I agreed to collect specimens and harvest the gills for DNA. I collected these small, red, russulas with a white

stem and very hot taste, before, during and after the foray. At the end of the season samples from 18 specimens were sent to Anna. All were from coniferous forests, but all had at least some birch in the vicinity. Their colour ranged from a very pale pink through a bright red to a dusker red. Some also showed tinges of violet, not evident on these photos. All were photographed, microscoped and dried. Andrus tells me that at least some of the pale ones, which we had called *R. aquosa*, had bigger spores, close to *R. sylvestris*, so we should find at least two species, possibly more.

Once Anna has results available, you shall get further updates.

Photo: Roger Smith



Foray Fotos

*Photos by : Andrus Voitk, Maria Voitk, Roger Smith,
Michael Burzynski, Marian Wissink*













Program

Friday, September 25

- 11:00 Mycoblitz at Sir Richard Squires Memorial Provincial Park (turn off Highway 430, eight kilometres north west of Deer Lake).
- 2:00 Leave to drive to Killdevil Camp.
- 4:00 Sign-in desk opens at Killdevil Camp.
- 6:00 *Reception/Supper* begins
- 7:30 *Words From the President*
- 8:00 Simultaneous Talks:
- *Mushrooms 101*, Faye Murrin.
 - *So Many Species, So Little Time: Fungal Biodiversity in the Great Smoky Mountains National Park*, Andrew Miller.

Saturday, September 26

- 8:00 Breakfast and announcements.
9:00 Foray teams leave for various trails.
12:00 Lunch on the trail.
1:00 Identifiers and databasers return to start identification and processing.
3:00 Foray teams return to sort, label, and identify specimens.
6:00 *Quidi Vidi QuuQup* (wild mushrooms and supper).
7:30 Short Talks:
 - *A New Look at an Old Genus: Lactarius*, Andrew Methven
 - *Mushroom Poetry*, Michel Savard
 - *Exploring Fluorescent Fungi*, Greg Thorn and Michele Piercey-Normore
9:00 Lichen night hike

Sunday, September 27

- 8:00 Breakfast.
8:45 Group Photograph.
9:00 Specimen Tables and Workshops:

| | | | | |
|-----------------------|--------------------------|--|---|--|
| 9:00 to 10:00 | Tables with Andrus | Watercolour with Glynn Bishop (max. 10) | Cooking Wild Mushrooms with Jason Nesbitt (max. 12) | Let it Rot Walk with Nils Hallenberg (max. 14) |
| 10:00 to 11:00 | Tables with Michele | | | |
| 11:00 to 12:00 | Tables with Andy Methven | Lichen Walk with Michele Piercey-Normore (max. 14) | Pick for the Pot with Michael and Anne (max. 12) | |
| 12:00 to 1:00 | Tables with Oluna | | | |

- 1:00 Lunch.
2:00 President's Thanks.
2:15 *Annual General Meeting*.
3:00 Foray 2015 Concludes.

Workshops and Walks

Sunday morning was devoted to workshops, both indoors and out: Let it Rot walk, Pick for the pot, Lichen walk, Watercolour sketching, Cooking Wild Mushrooms, and, of course, Tables sessions. To get a flavour of what they were about, here are some “Reports” by some of the leaders and participants.

Watercolour Workshop by Glynn Bishop

The watercolour workshop took place for the second year in the wonderful chapel that is tucked into the trees above the salt water inlet, and with the backdrop of Killdevil Mt. covered with a frosting of snow.

Participants were introduced to drawing and painting

outdoors with all its challenges. Watercolour techniques and drawing tips were blended with a mixture of wildlife awareness and seeing, as opposed to looking.

The workshop concentrated on identification illustration, and note taking combined with colour rendering.



Photos: Michael Burzynski

Pick for the Pot—Success at Last!

by Michael Burzynski and Anne Marceau

After our dismal luck during Foray 2014, I am happy to report that we had a good *Pick for the Pot* this year. We had an enthusiastic group of thirteen participants, and we found a good range of species despite it being a bit late in the year for some of the mushrooms.

This *Pick for the Pot* was conducted in three stages: First we got together in the tent and looked at specimens of edible mushrooms that had already been collected. We discussed identifying characteristics, how to collect and clean mushrooms in the field, and how to avoid poisonous species.

Then we walked the trails, meadow, and forest in the vicinity of Killdevil Camp collecting the edible species that we came across. We were about a week late for the best crop of Edible Bolete (*Boletus*

edulis), but there were still many that were good enough to pick. We also came across Orange Milkcap (*Lactarius deterrimus*), Field Waxcap (*Cuphophyllus pratensis*) and a large ring of Meadow Mushrooms (*Agaricus campestris*) in the field; In the forest we found Yellowlegs (*Craterellus tubaeformis*), Hedgehogs (*Hydnum repandum*), White Birch Bolete (*Leccinum holopus*), Gypsy (*Cortinarius caperatus*), and Bog Russula (*Russula paludosa*).

As the event drew to a close, we returned to the Camp and convened in the Meeting Hall to sort through all of the mushrooms that had been found. This was a chance to check that everyone had indeed collected only edible species, and that the mushrooms were still in good enough condition to eat (which gave us a chance to discuss the various insects, slugs, and mammals that also enjoy edible mushrooms). All in all, we had a successful and enjoyable event.



Photo: Michael Burzynski

Anne checking over mushrooms collected by participants, after the search.

Photo: Michael Burzynski

Lichen walk

Michele Piercey-Normore

The lichen walk took place along the first part of the Lomond trail. We had a small group of four people this year who had many questions about the biology of lichens and their natural products. As we walked, we discussed the interactions between the algal and fungal partners in lichens and how lichens use some of the natural products for their protection in nature. Our first encounter occurred just before we reached the trail - *Dibaeis baeomyces* was present as a pinkish-white carpet on the lawn where the grass had been removed and the soil was exposed. As we entered the forested trail, some of the beard lichens were hanging from tree branches, the yellow-green



beard lichens *Usnea* spp., *Alectoria sarmentosa*, and the brown horse-hair lichens *Bryoria* spp. Species differences among these lichens were discussed. We soon discovered several species of *Cladonia* either growing on the ground along the trail, at the base of trees, or on decaying logs such as *Cladonia ochrochlora*, *C. squamosa*, and *C. gracilis*. In some places the only evidence of lichen colonization were the tiny nondescript squamules of many *Cladonia* species but, while they completely covered the surface of some logs, they could not be differentiated at this stage of growth. We also discovered small patches of the leprose lichen (*Lepraria lobificans*),

which is now known as *L. finkii*. This lichen forms a powdery mass that covers wood or soil. It is called the leprose lichen because the powdery layer brushes off easily when physically disturbed by animals. We also found some of the pin or stubble lichens that commonly inhabit standing de-barked stumps and sometimes are found on the bark itself. These are often included with the lichens but many of them are non-lichenized fungi and some are parasitic on lichens growing within the wood. As the walk came to an end we mingled with other groups in the forest reflecting on our findings.

Night hike

Michele Piercey-Normore and Greg Thorn

This year two back-to-back offerings of a night hike attracted about 36 people. The hikes were designed to highlight the ability of lichens and other fungi to fluoresce when exposed to ultraviolet light. We first provided a short presentation to explain fluorescence, what the human eye sees, and why lichens and other fungi fluoresce. This is very different from bioluminescence, the biological production of visible light, as in *Panellus stipticus* or the luminous dinoflagellates such as *Noctiluca*. A bioluminescent organism uses an enzyme in its cells to act on the chemical luciferin, releasing light (usually greenish-white) in the process. In contrast, fluorescence in lichens or other fungi is a result of the secondary metabolites within the lichen or fungus that absorb the ultraviolet light and reflect visible light of different colors. In our demonstration, we provided the UV light to stimulate fluorescence, using hand-held UV flashlights. In the daytime, the lichens presumably fluoresce under the UV of sunlight, but their fluorescent output would be totally swamped by daylight - so that's why we had to do a night hike. Our first demonstration was the stipe bands on a basidiomycete, *Cortinarius armillatus*. (Next year, we need to remember to set aside a fresh specimen of the deadily, blue-fluorescent *Cortinarius rubellus*.) We also demonstrated fluorescent colors from the lichens on a large rock that was carried back from a previous field trip. The lichens on the rock included a number of map lichens (*Rhizocarpon* spp.),

Lecidea spp. and some small foliose lichens. These showed some yellows and purples when subjected to UV light. Then we walked to a big Red Maple tree in the front yard of Killdevil Camp. The trunk of the maple was colonized by many different lichens such as *Parmelia sulcata*, *Pertusaria macounii*, *Pertusaria amara*, *Ochrolechia androgyna*, and several *Lobaria* species. The *Pertusaria* spp. produced purple and pale red colors when exposed to the ultraviolet flashlight but the *Lobaria* and *Parmelia* species were not as striking and did not show fluorescence. The purple colors on the maple tree were from natural products such as xanthenes in the *Pertusaria* spp. and the pale yellows on the rock were from rhizocarpic acid in the map lichen (*Rhizocarpon* spp.). After several expressions of awe we hiked to the back of the camp where a lichen *Dibaeis baeomyces* covered the ground. This lichen fluoresced an orange and bluish-white color, which is due to the natural products baeomycesic acid and squamatic acid reflecting light in those wavelengths. The night hike was a success and may reveal some new fluorescent colors next year. This is definitely an idea you can try at home - but remember to wear UV-protective eye-wear. The orange, UV-absorbing plastic safety glasses supplied by Michael Burzynski made some of the fluorescence much more obvious, as well as providing protection to the wearer's retinas from UV damage.

Foraging for the Kitchen

By Milly Brown

Foray 2015 was my first (but likely not my last) adventure in learning about the gastronomic delights of mushroom foraging. The pinnacle of the experience was standing in a cozy circle with my fellow fungi hunters around Jason Nesbitt as he made the most wonderful Japanese-style soup with a mouth-watering collection of local wild mushrooms, both fresh and dried. Jason's cooking style was part art, part magic, and the resulting potion transported me to another place—a happy, comfortable and satisfied one. He trusts his instincts, and encouraged us to do so as well, measuring in dollops, splashes and other intuitive quantities. That works for me, but I did take detailed notes to help me create similar soups in the future....

Wild Mushroom Soup & Dumplings, Japanese-Style (a Jason Nesbitt recipe)

Ingredients

Chopped onion or shallot, celery, carrot, fresh ginger root
Whole small hot chili pepper
Dried lobster mushrooms and morels, soaked overnight & drained, reserving liquid for soup
Variety of fresh mushrooms (chanterelles, orange milk mushrooms, hedgehogs, slippery Jacks)
Garlic oil (see recipe below)
Tawny sherry
Fresh lime, juice & peel
Mirin
Seasoned rice wine
Soy sauce
Chicken stock
Wonton wrappers
Beaten egg
Panko crumbs
Wild chive puree
Noodles, cooked al dente
Garnish: chopped fresh coriander & green onions

Preparation

In advance, prepare

- Chicken stock by simmering chicken pieces in water with a little salt.
- Garlic oil by roasting whole garlic cloves for 10 minutes at 400 degrees, then storing them in a mason jar with 60%/40% olive oil/vegetable oil, sea salt, peppercorns & rosemary.
- Rehydrated dried mushrooms by soaking overnight in water.

Sauté chopped onion, celery, carrots, ginger in sesame oil with a splash of garlic oil for around 5 minutes, then start adding mushrooms, starting with the densest rehydrated ones, then moister fresh ones, slowly adding each and letting juices evaporate as you go. Slice a chili pepper once vertically and toss it in whole so that you can pull it out if the soup seems spicy enough. Add about 1/3 cup sherry & keep cooking until mostly evaporated. Grate a little lime peel into pot. Add a few tablespoons each of seasoned rice wine & mirin, 1 tbsp soy sauce, and the juice of half a lime. Keep reducing liquid until mixture is almost dry & caramelized. Turn off heat & rest it for a few minutes. You can use this mixture immediately over rice or noodles, or go on to make dumplings & soup...

Dumplings

Take about a cup of the sautéed mixture & pulse in a blender to chop finely, but NOT puree, & let cool. Add a beaten egg, ¼ cup Panko crumbs, & season to taste if needed with more soy sauce, sesame, chili and/or lime juice. Moisten the edges of a wonton wrapper with water, put a teaspoon of filling in the centre, fold over diagonally & pinch edges to seal. Lay out filled wontons in a single layer on a baking sheet dusted with cornstarch. If desired for advance preparation, you can blanch the filled dumplings briefly in chicken stock, then brush them with oil and layer with parchment paper until you are ready to add them to the soup.

Soup

To the remaining sautéed mix, add chicken stock, return to heat, & bring to a boil. Add reserved mushroom liquid & keep simmering. Season to taste with more soy sauce, sesame, chili and/or lime juice as desired. Add al dente noodles to soup. Turn up heat, add filled dumplings, and bring to boil. Turn off heat and serve garnished with fresh coriander & a few dots of pureed wild chives.



Photos: Michael Burzynski



Photo: Michael Bruzynski



**Let it Rot Walk
with Nils Hallenberg**

Photos: Marian Wissink



**Tables
with Andrus, Michele, Andy
and Oluna**



Photos: Marian Wissink

Photos: Michael Burzynski




Photo: Maria Voitek



Trail Finds



Photos: Michael Burzynski



*Sidewalk mushroom *Agaricus bitorquis* bursting through four inches of fresh asphalt Route 431.*

Trails

| | Gros Morne Mountain | Lomond River Trail | Stuckless Pond Trail | Stanleyville Trail | Green Gardens Trail (Wallace Brook) | Trout River Pond Trail |
|---|--|---|---|--|---|---|
| Terrain | Moist coniferous forest, heath barren, tuckamore, rock barren, rock knob | Rich fen, mixed forest, riverside, insect- and wind-damaged forest, meadow, estuary | Mixed forest, coniferous forest, lakeside, meadow | Mixed forest, insect- and wind- damaged forest, meadow, seacoast | Peridotite barrens, riverside, coniferous forest, seacoast | Mixed forest, lakeside, peridotite barrens, travertine seeps |
| Difficulty: 1-easy; 5-tough | Beginning of trail- 2 Entire trail- 5 | Beginning of trail- 2 Entire trail- 3 | Beginning of trail- 2 Entire trail- 3 | 3 | Beginning of trail- 2 Entire trail- 4 | Beginning of trail- 2 Entire trail- 3 |
| Productivity 1-low; 5-high | 5 | 4 | 4 | 3 | 3 | 3 |
| Points of Interest | Brook and rich forest understory | River and estuary; unusually diverse forest | Old logging trail, mature conifers | Route to the area's first lumber mill; limestone cliffs | Differences between vegetation of serpentine and normal soils | Glacial trough lake (a former fjord); views of the Tablelands |
| Length (km) See note below | 16 km (loop) | 6 km (one way) | 9.5 km (loop) | 4 km (return) | 9 km (return) | 14 km (return) |
| Start | GM Trail parking lot | LR Trail parking lot | LR Trail parking lot | Lomond Campground, Day-use-area parking lot | GG Trail, Wallace Brook parking lot | Trout River Pond Day-use-area parking lot |
| Leader | Jamie Graham | Helen Spencer | Marian Wissink | Jessica Butler | Anne Marceau | Maria Voitk |
| Databaser | Michael Burzynski | Tony Wright | Claudia Hanel | Roger Smith | Chris Deduke | Shane White |
| Mycologist | Nils Hallenberg | Andy Methven | Andy Miller | Greg Thorn | Oluna Ceska | Andrus Voitk |
| Other | Faye Murrin | | | Michele Piercey-Normore | | |
| Trail sign-up sheets will be posted during Registration on Friday. Please put your name on a trail that you would like to do. Note the limit of ten participants (including leaders) per trail. | | | | | | |

SPECIES LIST

DISTRIBUTION BY FORAY TRAIL

Non-lichenized fungi, basidiolichens and myxomycetes

*Andrus Voitk, Michael Burzynski, Tony Wright
with major input from the faculty*

Trails

KD = Killdevil grounds
LR = Lomond River Trail
SV = Stanleyville Trails
ST = Stuckless Pond Trail
TR = Trout River Pond Trail + Trout River Head Trail
GG = Green Gardens Trail (Long Pond entrance)
BH = Burnt Hill, Norris Point
LC = Lobster Cove Head lighthouse grounds & trails
GM = trail to Gros Morne Mountain
WB = Western Brook Pond Trails

BL = Big Level, GMNP

SRS = Sir Richard Squires Memorial Provincial Park,

O/U = other or unknown site

TOT column lists the number of collections of a species.

Remaining columns—same information for each trail.

NB: number of collections approximate indicator of abundance. Comparison of trails not valid, because all not sampled equally.

BLUE = surveyed by foray only

YELLOW = surveyed by Faculty only

GREEN = surveyed by both Faculty and Foray

Background: Spores of *Entomophthora muscae* on windowpane of Lobster Cove Head Lighthouse.

This species is the poster boy of our 2015 list—see discussion under **What do these data mean?** This is a very fascinating fungal species, and we very strongly encourage you to look it up on Tom Volk's Fungus of the Month series: <http://botit.botany.wisc.edu/toms_fungi/mar2000.html>.

Photo: Michael Burzynski

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|---|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Agaricus bitorquis</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Agaricus campestris</i> | 2 | 1 | | | | | | | 1 | | | | | |
| <i>Aleurodiscus amorphus</i> | 4 | | | 1 | | 1 | | | 1 | | | | 1 | |
| <i>Alloclavaria purpurea</i> | 1 | | | | | | | 1 | | | | | | |
| <i>Amanita bisporigera</i> | 4 | | 2 | 1 | 1 | | | | | | | | | |
| <i>Amanita flavoconia</i> | 3 | | | | | 1 | | 1 | | | | | 1 | |
| <i>Amanita fulva</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Amanita muscaria</i> var. <i>guessowii</i> | 12 | 1 | 2 | | | 1 | 2 | 2 | 1 | | | | 2 | 1 |
| <i>Amanita porphyria</i> | 4 | | 1 | | | | | | | 2 | | | 1 | |
| <i>Amphinema byssoides</i> | 9 | | | | 1 | 1 | | | 4 | | 1 | 1 | 1 | |
| <i>Ampulloclitocybe clavipes</i> | 5 | | | | | | | 2 | | 2 | 1 | | | |
| <i>Antrodia heteromorpha</i> | 5 | | 2 | 1 | | 1 | | | | 1 | | | | |
| <i>Antrodiella canadensis</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Antrodiella hoehnelii</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Armillaria ostoyae</i> | 3 | | 3 | | | | | | | | | | | |
| <i>Arrhenia acerosa</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Ascocoryne cylichnium</i> | 2 | | | | 1 | | | | | | | | 1 | |
| <i>Ascocoryne turficola</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Asterophora parasitica</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Athelicum</i> sp 1 | 1 | | | | | | | | 1 | | | | | |
| <i>Atheniella adonis</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Auricularia americana</i> | 2 | | 1 | 1 | | | | | | | | | | |
| <i>Austroboletus gracilis</i> | 4 | | 1 | | | 1 | 2 | | | | | | | |
| <i>Bankera violascens</i> | 2 | | | | | 1 | | | | 1 | | | | |
| <i>Basidioidendron caesiocinereum</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Basidioradulum radula</i> | 7 | | | | 1 | 3 | | | 2 | 1 | | | | |
| <i>Biscogniauxia repanda</i> | 1 | | | | 1 | | | | | | | | | |
| <i>Bisporella citrina</i> | 4 | | 1 | | | 1 | | | | | | | 2 | |
| <i>Bjerkandera adusta</i> | 2 | | 1 | | | | | | | | | | 1 | |
| <i>Bogbodia uda</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Boletus edulis</i> | 4 | | | | | 1 | | 2 | | | | | 1 | |
| <i>Boletus subtomentosus</i> f. <i>gracilis</i> | 2 | | 1 | | | | | | | | | | 1 | |
| <i>Botryobasidium subcoronatum</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Botryobasidium vagum</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Botryohypochnus isabellinus</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Bovista pila</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Brevicellicium exile</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Camarophylloopsis foetens</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Cantharellus</i> sp. NL | 6 | | | | | | | 5 | 1 | | | | | |
| <i>Catathelasma ventricosum</i> | 5 | | | | | | | 2 | 1 | | | | 2 | |
| <i>Ceraceomyces borealis</i> | 1 | | | | | 1 | | | | | | | | |

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|------------------------------------|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Ceraceomyces microsporus</i> | 2 | | | | | | | | | 1 | 1 | | | |
| <i>Chaetosphaeria longiseta</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Chlorociboria aeruginascens</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Chlorociboria aeruginosa</i> | 2 | | | | | 1 | | | | 1 | | | | |
| <i>Chondrostereum purpureum</i> | 1 | | | | 1 | | | | | | | | | |
| <i>Chrysomphalina chrysophylla</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Cinereomyces lindbladii</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Clavaria amoenoides</i> | 2 | | 1 | | | | | | | | | | 1 | |
| <i>Clavaria argillacea</i> | 2 | | | | | | | | | | | 2 | | |
| <i>Clavaria fusiformis</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Clavaria rubicundula</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Clavaria tenuipes</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Clavaria vermicularis</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Clavulina coralloides</i> | 12 | 1 | 3 | 1 | 1 | 2 | | | 1 | | 1 | 1 | 1 | |
| <i>Clitocybe candicans</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Clitocybe dealbata</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Clitocybula familia</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Clitopilus prunulus</i> | 1 | | | | | | | 1 | | | | | | |
| <i>Collybia tuberosa</i> | 4 | | | | | | 2 | 2 | | | | | | |
| <i>Coltricia perennis</i> | 3 | | | | | | | | | | | | 3 | |
| <i>Conferticium ochraceum</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Coniochaeta velutina</i> | 2 | | | | | | | | | | | | 2 | |
| <i>Coniophora arida</i> | 1 | | | | | | 1 | | | | | | | |
| <i>Coniophora puteana</i> | 2 | | | | | 2 | | | | | | | | |
| <i>Conocybe juniana</i> | 1 | | | | | | | | | | | | | 1 |
| <i>Coprinus comatus</i> | 4 | | | | | | | | | | | | 3 | 1 |
| <i>Cortinarius acutus</i> | 8 | | | 2 | 1 | 1 | | 2 | | | 2 | | | |
| <i>Cortinarius alboviolaceus</i> | 4 | | | | | 3 | | | | | | | 1 | |
| <i>Cortinarius anomalus</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Cortinarius armillatus</i> | 18 | 1 | 6 | 1 | 2 | 3 | | | | 1 | | | 4 | |
| <i>Cortinarius bivelus</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Cortinarius bolaris</i> | 2 | | | | | 2 | | | | | | | | |
| <i>Cortinarius camphoratus</i> | 10 | | 3 | 1 | | 1 | 1 | 1 | | 1 | | | 2 | |
| <i>Cortinarius caperatus</i> | 6 | | | | 1 | | 1 | 1 | | | | 1 | 2 | |
| <i>Cortinarius cinnamomeus</i> | 6 | | | | | 1 | | 1 | 1 | | 1 | | 2 | |
| <i>Cortinarius collinitus</i> | 3 | | | | | | | 1 | | | | 1 | 1 | |
| <i>Cortinarius croceus</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Cortinarius evernius</i> | 5 | | 1 | 1 | | | | | 1 | | | 2 | | |
| <i>Cortinarius flexipes</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Cortinarius fragrans</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Cortinarius gentilis</i> | 2 | | | | | | | 1 | | 1 | | | | |

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|---|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Cortinarius huronensis</i> | 4 | | | | | 1 | | | | 1 | | 1 | 1 | |
| <i>Cortinarius incognitus</i> | 2 | | | | | | | | | 1 | | | 1 | |
| <i>Cortinarius limonius</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Cortinarius malicorius</i> | 2 | | | | | | | | | 1 | | | | 1 |
| <i>Cortinarius pholideus</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Cortinarius rubellus</i> | 7 | | | | | 1 | | 3 | 2 | 1 | | | | |
| <i>Cortinarius stillatitius</i> | 3 | | | 1 | | 2 | | | | | | | | |
| <i>Cortinarius traganus</i> | 7 | 1 | | 1 | 2 | | | | | 1 | | | 2 | |
| <i>Cortinarius trivialis</i> | 3 | | | | | | | | | | | | 3 | |
| <i>Cortinarius uliginosus</i> | 2 | | | | | 1 | | | | | | 1 | | |
| <i>Cortinarius varius</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Cortinarius vibratilis</i> | 2 | | 1 | | | 1 | | | | | | | | |
| <i>Craterellus tubaeformis</i> | 19 | 1 | 3 | | 2 | 5 | 1 | 1 | 3 | 1 | 1 | 1 | | |
| <i>Crucibulum laeve</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Cudonia circinans</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Cudonia lutea</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Cuphophyllus borealis</i> | 3 | | 1 | | | 2 | | | | | | | | |
| <i>Cuphophyllus lacmus</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Cuphophyllus pratensis</i> | 6 | 1 | 1 | | | | | | | | | | 3 | 1 |
| <i>Cystoderma amianthinum</i> | 2 | | | | | 1 | | | | 1 | | | | |
| <i>Cytidia salicina</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Dacrymyces chrysospermus</i> | 13 | | 2 | | 2 | | 1 | | 3 | 2 | 1 | | 2 | |
| <i>Elaphocordyceps ophioglossoides</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Elaphomyces muricatus</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Encoelia furfuracea</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Entoloma cuniculorum</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Entoloma incanum</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Entoloma luridum</i> | 2 | 1 | | | | | 1 | | | | | | | |
| <i>Entoloma quadratum</i> | 4 | 1 | | 1 | | 2 | | | | | | | | |
| <i>Entomophthora muscae</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Exidiopsis calcea</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Flavophlebia sulphureoisabellina</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Fomes fomentarius</i> | 7 | | 2 | | | | 1 | | | | | | 4 | |
| <i>Fomitopsis ochracea</i> | 4 | | 1 | 1 | | | | | | 1 | | | 1 | |
| <i>Fomitopsis pinicola</i> | 6 | | 3 | 1 | 1 | | | | | | | | 1 | |
| <i>Fuligo septica</i> | 2 | | | | | | | 2 | | | | | | |
| <i>Galerina hybrida</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Galerina paludosa</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Galerina vittiformis</i> f. <i>bispora</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Ganoderma applanatum</i> | 2 | | | | | | | | | 1 | | | 1 | |
| <i>Gliophorus laetus</i> var. <i>flavus</i> | 1 | | 1 | | | | | | | | | | | |

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|--|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Gloeocystidiellum porosum</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Gloeophyllum protractum</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Gloeophyllum sepiarium</i> | 12 | | 1 | 1 | 1 | | 2 | 1 | | 1 | 2 | | 3 | |
| <i>Gloiothele citrina</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Gloioxanthomyces nitidus</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Gomphidius maculatus</i> | 1 | | | | | | 1 | | | | | | | |
| <i>Gomphus clavatus</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Gymnopilus junonius</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Gymnopilus picreus</i> | 2 | | | | | | | | | | 2 | | | |
| <i>Gymnopilus sapineus</i> | 4 | | 2 | 1 | 1 | | | | | | | | | |
| <i>Gymnopus dryophilus</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Gymnosporangium cornutum</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Hapalopilus nidulans</i> | 2 | | | | | | | | | | 1 | | 1 | |
| <i>Harrya chromapes</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Hebeloma candidipes</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Hebeloma geminatum</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Helminthosphaeria clavariarum</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Helvella lacunosa</i> | 6 | 2 | | | | | | | 2 | | | | 2 | |
| <i>Helvella macropus</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Hemimycena gracilis</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Henningsomyces candidus</i> | 2 | | | 1 | | | | | 1 | | | | | |
| <i>Hydnellum caeruleum</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Hydnellum multiceps</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Hydnellum peckii</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Hydnellum pineticola</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Hydnellum scrobiculatum</i> | 4 | | | 1 | | | | | | 3 | | | | |
| <i>Hydnum repandum</i> | 6 | | 1 | | | 2 | 1 | | | | | 1 | 1 | |
| <i>Hydnum repandum</i> var. <i>album</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Hydnum rufescens</i> | 3 | | 1 | | | | | | | 1 | | | 1 | |
| <i>Hydnum umbilicatum</i> | 14 | | 2 | | 2 | 5 | | | | 1 | | 2 | 1 | 1 |
| <i>Hygrocybe acutoconica</i> | 6 | 1 | | | | 1 | | | | | | 1 | 3 | |
| <i>Hygrocybe cantharellus</i> | 10 | | 2 | 3 | | 1 | | | | | 2 | | 2 | |
| <i>Hygrocybe conica</i> | 4 | 1 | | | | | | | | | | 1 | 1 | 1 |
| <i>Hygrocybe miniata</i> | 3 | | 1 | | | | | | | | | 1 | 1 | |
| <i>Hygrocybe phaeococcinea</i> | 3 | 1 | | | | | | | | | | | 1 | 1 |
| <i>Hygrocybe punicea</i> | 1 | | | | | | 1 | | | | | | | |
| <i>Hygrocybe tahquamenonensis</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Hygrocybe turunda</i> | 2 | | | | | 1 | | | | | 1 | | | |
| <i>Hygrophorus chrysodon</i> | 1 | | | | | | 1 | | | | | | | |
| <i>Hymenochaete fuliginosa</i> | 2 | | | | | | | | 1 | 1 | | | | |
| <i>Hyphoderma setigerum</i> | 4 | | | | | 1 | | | 1 | 1 | 1 | | | |

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|---|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Hyphoderma sibiricum</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Hyphodontia arguta</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Hyphodontia borealis</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Hyphodontia breviseta</i> | 3 | 1 | | | | 1 | | | | | 1 | | | |
| <i>Hypholoma capnoides</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Hypholoma elongatum</i> | 2 | | | | | | | | | 1 | | | 1 | |
| <i>Hypholoma marginatum</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Hypochnicium cremicolor</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Hypochnicium punctulatum</i> | 2 | | | | | 1 | | | | | | 1 | | |
| <i>Hypomyces chrysospermus</i> | 2 | | | | | 1 | | 1 | | | | | | |
| <i>Hypomyces hyalinus</i> | 3 | | | | | 1 | | 2 | | | | | | |
| <i>Hypomyces lateritius</i> | 3 | | | 2 | | | | | 1 | | | | | |
| <i>Hypoxydon fuscum</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Immersiella immersa</i> | 1 | | | | 1 | | | | | | | | | |
| <i>Inocybe geophylla</i> | 19 | 1 | | 2 | | 1 | 1 | 2 | 6 | | 1 | | 4 | 1 |
| <i>Inocybe lanuginosa</i> | 2 | | | 1 | | | | | 1 | | | | | |
| <i>Inocybe lilacina</i> | 4 | | | | | | | | | | | | 4 | |
| <i>Inocybe phaeodisca</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Inocybe splendens</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Inocybe stellatospora</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Inonotus obliquus</i> | 2 | | | | | | | | | | | | 2 | |
| <i>Jahnoporus hirtus</i> | 3 | | | 1 | | 1 | | | | | 1 | | | |
| <i>Kavinia alboviridis</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Kirschsteiniotelia aethiops</i> | 2 | | | | 2 | | | | | | | | | |
| <i>Kneiffiella abieticola</i> | 2 | | | | | | | | | | 1 | | 1 | |
| <i>Laccaria bicolor</i> | 8 | | | | 1 | | | 1 | | 2 | 2 | 1 | 1 | |
| <i>Laccaria laccata</i> | 3 | | | | | | | | | | | 2 | 1 | |
| <i>Lachnum sulphurellum</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Lachnum virgineum</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Lacrymaria lacrymabunda</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Lactarius affinis</i> | 3 | | | | | | | | | | | 1 | 2 | |
| <i>Lactarius aspideoides</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Lactarius camphoratus</i> | 21 | 1 | 5 | 2 | | 4 | | 2 | 4 | 1 | 1 | | 1 | |
| <i>Lactarius deceptivus</i> | 18 | 2 | 2 | 3 | | | | | | 2 | | | 9 | |
| <i>Lactarius deterrimus</i> | 21 | | 1 | | 1 | 5 | 2 | | 5 | 1 | 2 | | 3 | 1 |
| <i>Lactarius glyciosmus</i> | 10 | | | | | | | | | 4 | | 2 | 4 | |
| <i>Lactarius griseus</i> | 1 | | | | 1 | | | | | | | | | |
| <i>Lactarius helvus</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Lactarius hibbardiae</i> | 5 | | | | | | | | | | | | 5 | |
| <i>Lactarius hysginus</i> | 1 | | | | | | | | | | | | | 1 |
| <i>Lactarius lignyotus</i> var. <i>canadensis</i> | 8 | | 1 | | | 2 | | | 1 | 1 | | 1 | 1 | 1 |

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|--|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Lactarius lignyotus</i> var. <i>lignyotus</i> | 1 | | | | 1 | | | | | | | | | |
| <i>Lactarius lignyotus</i> var. <i>nigroviolascens</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Lactarius mucidus</i> | 2 | | | | | 1 | | | | 1 | | | | |
| <i>Lactarius oculatus</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Lactarius pubescens</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Lactarius repraesentaneus</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Lactarius resimus</i> | 3 | | | | | | | | | | | 1 | 2 | |
| <i>Lactarius scrobiculatus</i> var. <i>canadensis</i> | 2 | | | | | | 1 | | 1 | | | | | |
| <i>Lactarius sordidus</i> | 1 | | | | | | 1 | | | | | | | |
| <i>Lactarius tabidus</i> | 3 | | 1 | | | 1 | | | | 1 | | | | |
| <i>Lactarius thyinos</i> | 20 | | 3 | 2 | 2 | 2 | | 3 | 4 | 2 | | | 1 | 1 |
| <i>Lactarius torminosus</i> | 5 | 1 | | | | 1 | | | | | | | 2 | 1 |
| <i>Lactarius uvidus</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Lactarius vietus</i> | 2 | | 1 | | | | | | | | | | 1 | |
| <i>Lactarius vinaceorufescens</i> | 12 | | 2 | 1 | 1 | 3 | | 1 | | 1 | | | 3 | |
| <i>Lasiosphaeria ovina</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Leccinum holopus</i> | 18 | | 5 | 1 | 1 | 3 | 2 | | | 1 | | | 5 | |
| <i>Leccinum rotundifoliae</i> | 3 | | | | | | | | | | | 3 | | |
| <i>Leccinum scabrum</i> | 12 | | 2 | | 2 | | | 2 | | 1 | | | 4 | 1 |
| <i>Leccinum vulpinum</i> | 9 | | 1 | | | | | | 1 | 1 | | | 6 | |
| <i>Leotia lubrica</i> | 11 | | 1 | 1 | 1 | 2 | | | 4 | 2 | | | | |
| <i>Leotia viscosa</i> | 8 | | 3 | | 1 | 3 | | | | 1 | | | | |
| <i>Lepiota cristata</i> | 2 | 2 | | | | | | | | | | | | |
| <i>Leucoagaricus leucothites</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Lichenomphalia alpina</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Lichenomphalia hudsoniana</i> | 5 | | | | | | | | | | | 5 | | |
| <i>Lichenomphalia umbellifera</i> | 5 | | | | | | | | | | | 5 | | |
| <i>Lophium mytilinum</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Lycogala epidendrum</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Lycoperdon nigrescens</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Lycoperdon perlatum</i> | 7 | | | 1 | | | | 3 | 1 | 1 | | | 1 | |
| <i>Lycoperdon pyriforme</i> | 10 | | 1 | 1 | | | | | | | | 1 | 6 | 1 |
| <i>Lycoperdon umbrinum</i> | 2 | | | | | | 1 | | 1 | | | | | |
| <i>Lyophyllum decastes</i> | 2 | 2 | | | | | | | | | | | | |
| <i>Lyophyllum fumosum</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Marasmiellus perforans</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Melanoleuca brevipes</i> | 2 | | | | | | | 1 | | | | | 1 | |
| <i>Melanoleuca melaleuca</i> | 2 | 1 | | | | | | | | | | | 1 | |
| <i>Melanomma pulvis-pyrius</i> | 1 | | | | 1 | | | | | | | | | |
| <i>Melanospora caprina</i> | 1 | | | | | | | | | | | | 1 | |


| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|--|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Merismodes anomala</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Merismodes fasciculata</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Microglossum rufum</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Mollisia cinerea</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Mycena concolor</i> | 2 | | | | 2 | | | | | | | | | |
| <i>Mycena galericulata</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Mycena leptcephala</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Mycena niveipes</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Mycena pura</i> | 2 | | | | | | | | 2 | | | | | |
| <i>Neocudoniella albiceps</i> | 4 | | 1 | | 1 | | | | | 2 | | | | |
| <i>Neolecta irregularis</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Omphalina oreades</i> | 9 | | | | | | | | | | | 9 | | |
| <i>Onnia tomentosa</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Onygena equina</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Otidea onotica</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Panellus stipticus</i> | 2 | | | | | 1 | | | | | 1 | | | |
| <i>Paxillus involutus</i> | 8 | | | 1 | | | | | | 2 | 2 | | 3 | |
| <i>Peniophora cinerea</i> | 2 | | | | | | | | | | 2 | | | |
| <i>Peniophora erikssonii</i> | 2 | | | | | 2 | | | | | | | | |
| <i>Peniophora rufa</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Peniophorella pubera</i> | 2 | | | | | 1 | | | 1 | | | | | |
| <i>Peziza badia</i> | 2 | 1 | | 1 | | | | | | | | | | |
| <i>Peziza brunneoatra</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Peziza praetervisa</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Peziza vesiculosa</i> | 1 | | | | | | 1 | | | | | | | |
| <i>Phaeolus schweinitzii</i> | 4 | 2 | | | | | | | | | 1 | | | 1 |
| <i>Phellinus chrysoloma</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Phellinus ferreus</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Phellinus ferruginosus</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Phellinus laevigatus</i> | 2 | | | | | 1 | | | | | 1 | | | |
| <i>Phellinus viticola</i> | 2 | | | | | 1 | | 1 | | | | | | |
| <i>Phellodon niger</i> var. <i>niger</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Phellodon tomentosus</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Phlebia radiata</i> | 2 | | | | | | | | 1 | | 1 | | | |
| <i>Phlebia subserialis</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Phlebiella tulasnellodea</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Phlebiella vaga</i> | 2 | | | | | | | | 1 | 1 | | | | |
| <i>Pholiota astragalina</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Pholiota limonella</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Pholiota malicola</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Phyllotopsis nidulans</i> | 1 | | | | | | | | | | | | 1 | |

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|---|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Piloderma byssinum</i> | 2 | | | | | | | | | | | | 2 | |
| <i>Piptoporus betulinus</i> | 11 | | | 2 | 1 | | 1 | | | 1 | 1 | | 5 | |
| <i>Pleurocybella porrigens</i> | 3 | 1 | 2 | | | | | | | | | | | |
| <i>Plicatura nivea</i> | 2 | | | | | 1 | | | | | 1 | | | |
| <i>Plicaturopsis crispa</i> | 2 | | | | | 2 | | | | | | | | |
| <i>Pluteus plautus</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Pluteus podospileus</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Podophacidium xanthomelum</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Polyporus badius</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Polyporus varius</i> | 2 | | | 2 | | | | | | | | | | |
| <i>Postia caesia</i> | 4 | | | | 1 | 1 | | | 1 | | | | 1 | |
| <i>Postia fragilis</i> | 2 | | 2 | | | | | | | | | | | |
| <i>Postia ptychogaster</i> | 1 | | | | | | | 1 | | | | | | |
| <i>Postia stiptica</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Protostropharia alcis</i> | 6 | | 1 | | | | | | 1 | | | | 4 | |
| <i>Psathyrella conissans</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Psathyrella piluliformis</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Pseudohydnum gelatinosum</i> | 7 | | 1 | | | | | 1 | 5 | | | | | |
| <i>Puccinia poarum</i> | 2 | | | | | | | 1 | | | 1 | | | |
| <i>Radulomyces confluens</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Ramaria aurea</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Ramaria flaccida</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Ramaria flavicingula</i> | 3 | 1 | 1 | | 1 | | | | | | | | | |
| <i>Ramaria flavobrunnescens</i> | 2 | | | 2 | | | | | | | | | | |
| <i>Ramaria pallida</i> | 4 | | 1 | 2 | | | | | 1 | | | | | |
| <i>Ramaria suecica</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Ramariopsis crocea</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Resinicium bicolor</i> | 2 | | | | | | | | 1 | | 1 | | | |
| <i>Rhodocollybia maculata</i> | 1 | 1 | | | | | | | | | | | | |
| <i>Rhodocollybia maculata</i> var. <i>scorzoneria</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Rickenella fibula</i> | 6 | | | | | | 1 | | 2 | 2 | 1 | | | |
| <i>Roseodiscus subcarneus</i> | 2 | | 1 | 1 | | | | | | | | | | |
| <i>Russula adusta</i> | 5 | | | 1 | | | | | | 1 | | | 3 | |
| <i>Russula aeruginosa</i> | 1 | | | | | | | 1 | | | | | | |
| <i>Russula aquosa</i> | 2 | | 1 | | | | | | 1 | | | | | |
| <i>Russula brevipes</i> | 4 | | | | | 1 | | 2 | | 1 | | | | |
| <i>Russula claroflava</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Russula compacta</i> | 1 | | | | | | 1 | | | | | | | |
| <i>Russula decolorans</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Russula densifolia</i> | 2 | | | | | 2 | | | | | | | | |

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|------------------------------------|-----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|
| <i>Russula griseascens</i> | 8 | | 1 | | 1 | 1 | | | | 2 | 1 | | 2 | |
| <i>Russula laurocerasi</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Russula paludosa</i> | 7 | | 1 | | | 1 | | | 1 | 1 | | 2 | 1 | |
| <i>Russula peckii</i> | 13 | | 2 | 2 | 1 | 1 | | 1 | 2 | | | 1 | 2 | 1 |
| <i>Russula puellaris</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Russula variata</i> | 3 | | 1 | | | | | | | | | | 1 | 1 |
| <i>Sarcodon imbricatus</i> | 2 | | | 1 | | | | | | 1 | | | | |
| <i>Sarcodon stereosarcinon</i> | 4 | 2 | 1 | 1 | | | | | | | | | | |
| <i>Scopuloides rimosa</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Scytinostroma jacksonii</i> | 2 | | | | | | | 2 | | | | | | |
| <i>Sebacina epigaea</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Sebacina incrustans</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Sidera lenis</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Sistotrema confluens</i> | 1 | | | | | | | | | 1 | | | | |
| <i>Skeletocutis subincarnata</i> | 4 | | | | | 1 | | | | 1 | | | 2 | |
| <i>Sphaerobolus stellatus</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Sporormiella dubia</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Stereum hirsutum</i> | 2 | | | | | | | | | | 2 | | | |
| <i>Stereum rugosum</i> | 2 | | | | | 1 | | | | 1 | | | | |
| <i>Stereum sanguinolentum</i> | 7 | | 1 | | 1 | 1 | | | 2 | | 1 | | 1 | |
| <i>Suillus cavipes</i> | 10 | | | | | 1 | 3 | | | | | | 6 | |
| <i>Suillus clintonianus</i> | 9 | | | | 2 | | 1 | | | | | | 5 | 1 |
| <i>Suillus glandulosus</i> | 2 | | | | | 1 | | | | | | | 1 | |
| <i>Suillus grevillei</i> | 9 | | | | 2 | | 4 | | | | 1 | | 2 | |
| <i>Suillus grisellus</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Suillus paluster</i> | 2 | | | | | 1 | 1 | | | | | | | |
| <i>Suillus serotinus</i> | 4 | | | | 2 | | 1 | | | | | | | 1 |
| <i>Suillus spectabilis</i> | 5 | | | | | | 2 | | | | 2 | | 1 | |
| <i>Suillus spraguei</i> | 1 | | | | 1 | | | | | | | | | |
| <i>Taphrina robinsoniana</i> | 3 | | | | | 2 | | | | | 1 | | | |
| <i>Thelephora palmata</i> | 3 | | | | | | | 2 | 1 | | | | | |
| <i>Tomentella umbrinospora</i> | 2 | | | | | | | | 1 | | 1 | | | |
| <i>Tomentellopsis echinospora</i> | 1 | | | | | | | | | | | 1 | | |
| <i>Trametes ochracea</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Trametes versicolor</i> | 3 | | | | | 3 | | | | | | | | |
| <i>Trechispora confinis</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Trechispora farinacea</i> | 4 | | | | | | | | 1 | 1 | 1 | 1 | | |
| <i>Trechispora mollusca</i> | 3 | | | | | | 1 | | | | | 1 | 1 | |
| <i>Trechispora subsphaerospora</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Tremella foliacea</i> | 2 | | 1 | | 1 | | | | | | | | | |
| <i>Tremella mesenterica</i> | 1 | | | | | 1 | | | | | | | | |

| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|-------------------------------------|-------------|-----------|------------|-----------|-----------|------------|-----------|-----------|------------|-----------|-----------|-----------|------------|-----------|
| <i>Tremiscus helvelloides</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Trichaptum abietinum</i> | 4 | | | | 1 | | 1 | | | 1 | | | 1 | |
| <i>Trichaptum fuscoviolaceum</i> | 2 | | | | | | | | | | 2 | | | |
| <i>Trichaptum laricinum</i> | 4 | | 1 | 1 | 1 | | | | | 1 | | | | |
| <i>Trichaptum subchartaceum</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Trichodelitschia bisporula</i> | 1 | | 1 | | | | | | | | | | | |
| <i>Tricholoma apium</i> | 1 | | | 1 | | | | | | | | | | |
| <i>Tricholoma aurantium</i> | 1 | | | | 1 | | | | | | | | | |
| <i>Tricholoma pessundatum</i> | 1 | | | | | | | 1 | | | | | | |
| <i>Tricholoma saponaceum</i> | 2 | | | | | | | 2 | | | | | | |
| <i>Tricholoma subluteum</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Tricholoma subsejunctum</i> | 2 | | | | | | | 1 | | | | | 1 | |
| <i>Tricholoma transmutans</i> | 6 | | | | | 2 | | | 1 | 2 | | | | 1 |
| <i>Tricholoma vaccinum</i> | 5 | | | | | 3 | | | | 1 | | | 1 | |
| <i>Tricholoma virgatum</i> | 1 | | | | | | | 1 | | | | | | |
| <i>Tricholomopsis sulphureoides</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Tubaria furfuracea</i> | 2 | | | | | | | | | | | | 2 | |
| <i>Tubulicrinis glebulosus</i> | 1 | | | | | | | | | | 1 | | | |
| <i>Tubulicrinis subulatus</i> | 3 | | | | | | | | 1 | | 2 | | | |
| <i>Turbinellus floccosus</i> | 5 | | | | 1 | 2 | | 2 | | | | | | |
| <i>Tylopilus felleus</i> | 1 | | | | | | | | | | | | 1 | |
| <i>Tyromyces chioneus</i> | 3 | | | | | | 1 | | | 1 | | | 1 | |
| <i>Uredinopsis americana</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Xanthoporia radiata</i> | 2 | | | | | | | | | 2 | | | | |
| <i>Xerocomus badius</i> | 2 | | | | | | | | | | | 1 | 1 | |
| <i>Xeromphalina</i> sp. | 5 | | 1 | 2 | 1 | 1 | | | | | | | | |
| <i>Xylodon asperus</i> | 5 | | | | | 1 | | | | | 1 | 1 | 2 | |
| <i>Xylodon borealis</i> | 1 | | | | | 1 | | | | | | | | |
| <i>Xylodon nespori</i> | 1 | | | | | | | | 1 | | | | | |
| <i>Zignoëlla ovoidea</i> | 1 | | | | | | | | | | | | 1 | |
| Total collections | 1159 | 48 | 116 | 72 | 60 | 149 | 46 | 69 | 109 | 92 | 76 | 66 | 234 | 22 |
| Total taxa | 397 | 42 | 75 | 57 | 48 | 102 | 34 | 43 | 73 | 74 | 64 | 42 | 132 | 22 |
| TAXON | TOT | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |

| | KD | LR | SV | ST | TR | GG | BH | LC | GM | WB | BL | SRS | O/U |
|-------------|----|-----|----|----|-----|----|----|-----|----|----|----|-----|-----|
| Taxa | 42 | 75 | 57 | 48 | 102 | 34 | 43 | 73 | 74 | 64 | 42 | 132 | 22 |
| Collections | 48 | 116 | 72 | 60 | 149 | 46 | 69 | 109 | 92 | 76 | 66 | 234 | 22 |



Dead Cluster Fly infected with Entomophthora muscae.



What do these data mean?

Andrus Voitk



Lobster Cove Head Lighthouse

Photo: Roger Smith

The full-page photos introducing the list and this interpretation are purposely selected for their aptness. The first shows, somewhat vaguely, spores of an entity whose existence has escaped the awareness of most people, including those with a keen interest in fungi. It is the poster boy of our new species for 2015, a change that has been happening for the last few years. The 2013 data interpretation said, "The list of new species ... contains relatively few of the large fleshy mushrooms that we think of as "normal" mushrooms." In 2014 the same analysis stated, "After 10 years of surveying various regions of the province with both small and large forays, we may have

uncovered most of the macrofungi in the province." Table 1 compares our new species to the same number of species arranged in decreasing abundance of collections. How many names in the column of new species do you recognize? Or, better, how many of the names convey even the vaguest of meaning to you? Now, how many names in the column of common species do you recognize? Even if this was your first foray, some of those names will be familiar to you, and you will recognize several more, even if you may not be entirely certain you actually know their owners. In other words, although our species list continues to rise (Figure 1), new

species now come from outside the experience of most of us.

Reviewing the new species reveals that a large proportion is contributed by subspecialists. Over 30 new species were corticioid fungi identified by Nils Hallenberg. The contribution of Nils to our new species would have been more than twice as big, had he not attended our foray in 2012: another 30-40 polypores and corticioids that he identified this time, he first identified in 2012. And in 2012 we noted that had Leif Ryvarden not been here the year before, Nils' contribution at that time would have been much larger. Another 10 new species were pyrenomycetes (the "little

Table 1. Comparison of NEW species and COMMON species, Foray 2015

| NEW species | Common species in decreasing order of commonness | NEW species | Common species in decreasing order of commonness |
|---|---|------------------------------------|---|
| <i>Antrodiella canadensis</i> | <i>Lactarius camphoratus</i> | <i>Lacrymaria lacrymabunda</i> | <i>Lycoperdon perlatum</i> |
| <i>Antrodiella hoehnelii</i> | <i>Lactarius deterrimus</i> | <i>Lasiosphaeria ovina</i> | <i>Pseudohydnum gelatinosum</i> |
| <i>Athelium sp 1</i> | <i>Lactarius thyinos</i> | <i>Lophium mytilinum</i> | <i>Russula paludosa</i> |
| <i>Biscogniauxia repanda</i> | <i>Craterellus tubaeformis</i> | <i>Melanoleuca brevipes</i> | <i>Stereum sanguinolentum</i> |
| <i>Botryohypochnus isabellinus</i> | <i>Inocybe geophylla</i> | <i>Melanomma pulvis-pyrius</i> | <i>Cantharellus sp. NL</i> |
| <i>Brevicellicium exile</i> | <i>Cortinarius armillatus</i> | <i>Melanospora caprina</i> | <i>Cortinarius caperatus</i> |
| <i>Camarophyllopsis foetens</i> | <i>Lactarius deceptivus</i> | <i>Mycena niveipes</i> | <i>Cortinarius cinnamomeus</i> |
| <i>Chaetosphaeria longiseta</i> | <i>Leccinum holopus</i> | <i>Neocudoniella albiceps</i> | <i>Cuphophyllus pratensis</i> |
| <i>Chrysomphalina chrysophylla</i> | <i>Hydnum umbilicatum</i> | <i>Omphalina oreades</i> | <i>Fomitopsis pinicola</i> |
| <i>Clavaria amoenoides</i> | <i>Dacrymyces chrysospermus</i> | <i>Peziza brunneoatra</i> | <i>Helvella lacunosa</i> |
| <i>Clavaria fusiformis</i> | <i>Leccinum scabrum</i> | <i>Peziza vesiculosa</i> | <i>Hydnum repandum</i> |
| <i>Clavaria rubicundula</i> | <i>Russula peckii</i> | <i>Phlebia radiata</i> | <i>Hygrocybe acutoconica</i> |
| <i>Clavaria tenuipes</i> | <i>Amanita muscaria var. guessowii</i> | <i>Phlebia subserialis</i> | <i>Rickenella fibula</i> |
| <i>Conferticium ochraceum</i> | <i>Clavulina coralloides</i> | <i>Phlebiella tulasnellodea</i> | <i>Protostropharia alcis</i> |
| <i>Coniochaeta velutina</i> | <i>Gloeophyllum sepiarium</i> | <i>Pholiota malicola</i> | <i>Tricholoma transmutans</i> |
| <i>Conocybe juniana</i> | <i>Lactarius vinaceorufescens</i> | <i>Piloderma byssinum</i> | <i>Ampulloclitocybe clavipes</i> |
| <i>Cudonia lutea</i> | <i>Leotia lubrica</i> | <i>Pluteus plautus</i> | <i>Antrodia heteromorpha</i> |
| <i>Entoloma cuniculorum</i> | <i>Piptoporus betulinus</i> | <i>Pluteus podospileus</i> | <i>Catathelasma ventricosum</i> |
| <i>Entomophthora muscae</i> | <i>Cortinarius camphoratus</i> | <i>Ramaria aurea</i> | <i>Cortinarius evernius</i> |
| <i>Exidiopsis calcea</i> | <i>Hygrocybe cantharellus</i> | <i>Ramaria flavicingula</i> | <i>Lactarius hibbardiae</i> |
| <i>Flavophlebia sulphureoisabellina</i> | <i>Lactarius glyciosmus</i> | <i>Ramaria suecica</i> | <i>Lactarius torminosus</i> |
| <i>Galerina hybrida</i> | <i>Lycoperdon pyriforme</i> | <i>Roseodiscus subcarneus</i> | <i>Lichenomphalia hudsoniana</i> |
| <i>Galerina vittiformis f. bispora</i> | <i>Suillus cavipes</i> | <i>Russula aeruginosa</i> | <i>Lichenomphalia umbellifera</i> |
| <i>Gliophorus laetus var. flavus</i> | <i>Amphinema byssoides</i> | <i>Scopuloides rimosa</i> | <i>Russula adusta</i> |
| <i>Hebeloma candidipes</i> | <i>Leccinum vulpinum</i> | <i>Scytinostroma jacksonii</i> | <i>Suillus spectabilis</i> |
| <i>Hydnum repandum var. album</i> | <i>Omphalina oreades</i> | <i>Sebacina epigaea</i> | <i>Tricholoma vaccinum</i> |
| <i>Hygrocybe tahquamenonensis</i> | <i>Suillus clintonianus</i> | <i>Sebacina incrustans</i> | <i>Turbinellus floccosus</i> |
| <i>Hypodontia borealis</i> | <i>Suillus grevillei</i> | <i>Sporormiella dubia</i> | <i>Xeromphalina sp.</i> |
| <i>Hypochnicium cremicolor</i> | <i>Cortinarius acutus</i> | <i>Tomentella umbrinospora</i> | <i>Xylodon asperus</i> |
| <i>Hypochnicium punctulatum</i> | <i>Laccaria bicolor</i> | <i>Tomentellopsis echinospora</i> | <i>Aleurodiscus amorphus</i> |
| <i>Immersiella immersa</i> | <i>Lactarius lignyotus var. canadensis</i> | <i>Trechispora mollusca</i> | <i>Amanita bisporigera</i> |
| <i>Inocybe phaeodisca</i> | <i>Leotia viscosa</i> | <i>Trechispora subsphaerospora</i> | <i>Amanita porphyria</i> |
| <i>Inocybe splendens</i> | <i>Paxillus involutus</i> | <i>Trichodelitschia bisporula</i> | <i>Bisporella citrina</i> |
| <i>Kavinia alboboviridis</i> | <i>Russula griseascens</i> | <i>Tricholoma subluteum</i> | <i>Boletus edulis</i> |
| <i>Kirschsteiniothelia aethiops</i> | <i>Basidioradulum radula</i> | <i>Tubulicrinis subulatus</i> | <i>Collybia tuberosa</i> |
| <i>Kneiffiella abieticola</i> | <i>Cortinarius rubellus</i> | <i>Xylodon borealis</i> | <i>Coprinus comatus</i> |
| <i>Lachnum sulphurellum</i> | <i>Cortinarius traganus</i> | <i>Xylodon nespori</i> | <i>Cortinarius alboboviolaceus</i> |
| <i>Lachnum virgineum</i> | <i>Fomes fomentarius</i> | <i>Zignoëlla ovoidea</i> | <i>Cortinarius huronensis</i> |




Figure 1. Orange line is the cumulative new species list over the years of forays. Green bars represent the number of species collected per foray (not all are comparable). The steady rise of the orange line, with no sign of levelling off, suggests that at nearly 1,500 species we are nowhere near having recovered most of our species. However, as the text makes clear, most of the obvious macrofungi may be identified. Additional species will be recovered among less familiar groups of fungi, or after phylogenetic study of large common groups. The number of “typical mushroom” species in the province may be less than 2,000.

black dots’’) contributed by Andy Miller. In both cases, most of these collections were also made by the same identifiers. Thus, clearly there are large groups of fungi that are not familiar to us. This is not to say that our generalist identifiers could not have identified these species, but first, they are so unfamiliar to us that we do not even collect them, and second, even were we to bring some in, our identifiers are so busy with more collections than they can handle, that additional material would likely have to be left fallow.

Now, you may notice that you recognize several members of the genera *Clavaria* and *Ramaria* in the list of new species. Indeed, this is so, but review suggests that some old species may have had different names applied this time. We shall encounter this again—new names, but not necessarily new species. Since then several ramarias have been sequenced, but **to sort them out is a thesis project. Any takers?**

The above aside, the new species list does not include any species from the large, common genera. We have 152 species of *Cortinarius*

on our cumulative list and identified 26 species this year; but none new. Similarly, look at *Amanita*, *Entoloma*, *Lactarius*, *Russula* or *Tricholoma*. Some of these genera do have a single representative on the new list, but again in several cases it is a matter of a mushroom that we have identified by another name in the past—new name, not necessarily new species. While we try to eliminate synonymy, when we cannot be certain, we prefer to let an additional name stand until more detailed investigations settle the issue. Where our identifications have been imprecise, several names may have been used for the same entity. In addition to the clavarias, for examples, see Figure 2, or the *Russula griseascens* survey update elsewhere in this issue.

The most interesting case is that of *Lactarius*, a large genus where we had a specialist present. Surely that should have brought out new species? Well, apparently not. *Lactarius* is far more familiar to us than *Kirschsteiniothelia*, *Scytinostroma* or *Zignöella*, so we are likely to have collected and identified far

more species of *Lactarius* over the years than of the other three. Does this mean that we have identified all our species of *Lactarius*, and by extension, other common, fleshy fungi? Not likely. But our ability to be more accurate with morphologic methodology, backed by available classical descriptions, is limited. These large genera and the exact species of them that we have here, will require phylogenetic analysis, where possible coupled to type studies. Andy Methven, our *Lactarius* specialist this year, knows this. For example, he signed off several species as *Lactarius deterrimus*, even though he knows that this European species may not exist in North America at all. He qualified one such identification with “sensu Amer. auct.,” which means according to the current interpretation of most American authors. Because of this, he agreed to help us try to sort out at least this group of *Lactarius*, so that in some future year we can use correct names for the group currently flying under the incorrect name, *Lactarius deterrimus*. Look for it in the future. Meanwhile, we continue writing



Photo: Roger Smith

Figure 2. *Tricholoma subluteum*, an example of where study of a common group could improve our species list. In past years, tricholomas like this have been variously identified as *T. fumosoluteum* or even *T. intermedium*. With less peaked caps

and entirely yellow gills, the *T. equestre* complex comes into play, and with more peaked caps, taller stems and more olive hues in addition to the yellow, the *T. subsejunctum* complex may cause confusion. In pine forests, *T. davisiae* needs to be considered—

very similar pine associates with more acute caps. Phylogenetic study of these groups should tell us which species we share with Europe, which are distinct on this continent, how many of these are found in NL, and how to identify them.

names currently in use, even if we know they are wrong, simply because we do not have better names to offer.

What does all this mean to you? Well, if your main interest is to spend an enjoyable week-end in the woods with likeminded companions, to learn a few things about the mushrooms with which we share this earth, to recognize a few species that may widen your dining pleasure, and then return home a little more knowledgeable and a little more able to appreciate the value of nature—no change. The foray will always try to offer you that because that is the main purpose of FNL.

If your aim is to identify most of the “traditional mushrooms” in the province, the task is not hopeless. Although we may have 6-8,000 fungal species here, “normal” mushroom species are probably under 2,000, a manageable number,

not all that different from the number of vascular plant species.

If you have other more scientific goals, then these data tell you that future efforts should be aimed in two directions.

1. Continue inviting subspecialists to add depth to groups otherwise rarely explored. Monitor the results to know when you are getting there. For example, if you know that there are over 10,000 species of pyrenomycetes, then the 20 or so on our list is unlikely to have exhausted the supply.
2. Work with investigators interested in our large groups, to study their phylogeny in our province. At this stage, we will not identify too many new species morphologically, but may discover that the species concepts we have used may be incorrect or may cover several closely similar species. Because we have a good amount of experience, as well as

voucher specimens and photographs, knowing genetic clades may allow us to look back and identify characters that help separate members of the clades from each other. Where possible, tying them to type material will help identify our species with certainty. *Agaricus*, *Amanita*, *Cortinarius*, *Lactarius*, *Ramaria*, *Russula*, and *Tricholoma* are a few of the bigger common genera that would benefit from such treatment.

The beacon of the Lobster Cove Head lighthouse provided sailors with information when to alter course to get to home port safely. Analysis of our data should provide similar information for changes in direction to learn more of our fungi. If this is not your main interest, but, like most of us, you enjoy the foray experience for itself, then surely it will be that much more rewarding to know that your participation created a helpful slipstream for the scientific ship to sail in your wake.

SPECIES LIST AND DISTRIBUTION BY FORAY TRAIL - LICHENS

Chris Deduke and Michele Piercey-Normore

| Species | Sir Richard Squires | Stanleyville | Western Brook Pond | Big Level | #2 Lomond River | Burnt Hill | Trout River Trail | Green Garden | Lobster Cove Head | Stuckless Pond | Kildevil Campground |
|---|---------------------|--------------|--------------------|-----------|-----------------|------------|-------------------|--------------|-------------------|----------------|---------------------|
| <i>Alectoria sarmentosa</i> | √ | √ | √ | | | | | | | | |
| <i>Amandinea punctata</i> | | | | | | | | | | | |
| <i>Arctoparmelia centrifuga</i> | | | | √ | | | | | | | |
| <i>Arctoparmelia incurva</i> | | | | √ | | | | | | | |
| <i>Athallia holocarpa</i> | | √ | | | | | | | | | |
| <i>Baeomyces rufus</i> | | √ | | √ | | | | | | | |
| <i>Bryoria trichodes</i> | √ | | √ | | | | | | | | |
| <i>Buellia stillingiana</i> (?) | √ | | | | | | | | | | |
| <i>Calicium abietinum</i> | | | | | √ | | | | | | |
| <i>Calicium</i> sp. | | √ | | | | | | | | | |
| <i>Carbonea vorticosa</i> | | | | | | √ | | | | | |
| <i>Cetraria aculeata</i> | | | | √ | | | | | | | |
| <i>Cetraria ericetorum</i> | | | | √ | | | | | | | |
| <i>Cetraria islandica</i> | | | √ | √ | | | | | | | |
| <i>Chaenotheca brunneola</i> | | | | | √ | | √ | | | | |
| <i>Chaenotheca furfuracea</i> | | | | | √ | | | | | | |
| <i>Cladonia acuminata</i> | | | | √ | | | | | | | |
| <i>Cladonia amaurocraea</i> | | | | √ | | | | | | | |
| <i>Cladonia arbuscula</i> | | | √ | √ | | | | √ | | | |
| <i>Cladonia arbuscula</i> ssp. <i>mitis</i> | | | √ | √ | | | | | | | |
| <i>Cladonia bellidiflora</i> | | | | √ | | | | | | | |
| <i>Cladonia borealis</i> | | | | √ | | | | √ | | | |
| <i>Cladonia boryi</i> | | | √ | √ | | | | √ | | | |
| <i>Cladonia caespiticia</i> | | √ | √ | | | | √ | | √ | | |
| <i>Cladonia cenotea</i> | √ | √ | √ | | | | | √ | √ | | |
| <i>Cladonia chlorophaea</i> | √ | √ | √ | | | | | | √ | | |
| <i>Cladonia coccifera</i> | | | | √ | | | | | | | |
| <i>Cladonia crispata</i> | √ | √ | √ | √ | | | | √ | | | |

| Species | Sir Richard Squires | Stanleyville | Western Brook Pond | Big Level | #2 Lomond River | Burnt Hill | Trout River Trail | Green Garden | Lobster Cove Head | Stuckless Pond | Kildevil Campground |
|---|---------------------|--------------|--------------------|-----------|-----------------|------------|-------------------|--------------|-------------------|----------------|---------------------|
| <i>Cladonia cristatella</i> | | | | ✓ | | | | ✓ | | | |
| <i>Cladonia deformis</i> | ✓ | ✓ | | | | | | | | | |
| <i>Cladonia digitata</i> | ✓ | ✓ | ✓ | | | | | | | | |
| <i>Cladonia fimbriata</i> | | | ✓ | ✓ | | | | | | | |
| <i>Cladonia gracilis</i> | | ✓ | | | | | | | | | |
| <i>Cladonia gracilis</i> ssp. <i>gracilis</i> | | | | ✓ | | | | | | | |
| <i>Cladonia macilenta</i> | ✓ | | ✓ | | ✓ | | | | | | |
| <i>Cladonia macrophylla</i> | | | | ✓ | | | | | | | |
| <i>Cladonia maxima</i> | | | | ✓ | | | | ✓ | | | |
| <i>Cladonia merochlorophaea</i> | | ✓ | | | | | | | | | |
| <i>Cladonia multiformis</i> | | | | | | | | ✓ | | | |
| <i>Cladonia ochrochlora</i> | ✓ | ✓ | ✓ | | ✓ | | | | ✓ | | |
| <i>Cladonia pleurota</i> | ✓ | | ✓ | ✓ | ✓ | | | ✓ | ✓ | | |
| <i>Cladonia pyxidata</i> | | ✓ | | | | | | ✓ | | | |
| <i>Cladonia rangiferina</i> | ✓ | | | ✓ | | ✓ | | ✓ | | | |
| <i>Cladonia rei</i> | ✓ | ✓ | ✓ | | | | | | | | |
| <i>Cladonia scabriuscula</i> | | ✓ | | | | | | | ✓ | | |
| <i>Cladonia squamosa</i> | ✓ | | ✓ | ✓ | | | | | ✓ | | |
| <i>Cladonia stellaris</i> | | | ✓ | ✓ | | ✓ | | ✓ | | | |
| <i>Cladonia stygia</i> | ✓ | | ✓ | ✓ | | | | ✓ | | | |
| <i>Cladonia subulata</i> | | ✓ | | | | | | | | | |
| <i>Cladonia turgida</i> | | | | | | | | ✓ | | | |
| <i>Cladonia uncialis</i> | | | ✓ | ✓ | | | | | | | |
| <i>Cladonia verticillata</i> | ✓ | | | ✓ | | | | | | | |
| <i>Cladonia wainioi</i> | | | | ✓ | | | | | | | |
| <i>Evernia mesomorpha</i> | | | | | | | | ✓ | | | |
| <i>Flavocetraria cucullata</i> | | | | ✓ | | | | | | | |
| <i>Flavocetraria nivalis</i> | | | | ✓ | | | | | | | |
| <i>Graphis scripta</i> | ✓ | ✓ | | | | | ✓ | | | | |
| <i>Hypogymnia incurvoides</i> | ✓ | ✓ | | | | | | | | | |
| <i>Hypogymnia krogiae</i> | | | ✓ | | | | | | | | |
| <i>Hypogymnia physodes</i> | ✓ | ✓ | ✓ | ✓ | ✓ | | | | ✓ | | |
| <i>Hypogymnia tubulosa</i> | ✓ | ✓ | ✓ | ✓ | ✓ | | ✓ | | | | |

| Species | Sir Richard Squires | Stanleyville | Western Brook Pond | Big Level | #2 Lomond River | Burnt Hill | Trout River Trail | Green Garden | Lobster Cove Head | Stuckless Pond | Kildevil Campground |
|---|---------------------|--------------|--------------------|-----------|-----------------|------------|-------------------|--------------|-------------------|----------------|---------------------|
| <i>Icmadophila ericetorum</i> | ✓ | | | | | | | | | | |
| <i>Lecanora polytropa</i> | | | | ✓ | | | | | | | |
| <i>Lecanora symmicta</i> | | | ✓ | | | | | | | | |
| <i>Lecidea albofuscescens</i> (?) | ✓ | | | | | | | | | | |
| <i>Lepraria finkii</i> | ✓ | | ✓ | | | ✓ | | | ✓ | | |
| <i>Lobaria pulmonaria</i> | ✓ | | ✓ | | ✓ | | | | ✓ | ✓ | |
| <i>Lobaria quercizans</i> | | | ✓ | | | | | | | | |
| <i>Lobaria scrobiculata</i> | | | ✓ | | | | | | ✓ | | |
| <i>Loxospora elatina</i> | ✓ | | ✓ | | ✓ | ✓ | | | ✓ | | |
| <i>Loxospora ochrophaea</i> | ✓ | ✓ | ✓ | ✓ | ✓ | | | | | | |
| <i>Melanelia hepatizon</i> | | | | ✓ | | | | | | | |
| <i>Melanelia stygia</i> | | | | ✓ | | | | | | | |
| <i>Montanelia panniformis</i> | | | | ✓ | | | | | | | |
| <i>Mycoblastus sanguinarius</i> | ✓ | | | ✓ | | | | ✓ | | | |
| <i>Mycocalicium subtile</i> (?) | | ✓ | | | | | | | | | |
| <i>Nephroma arcticum</i> | | | | ✓ | | | | | | | |
| <i>Ochrolechia androgyna</i> | | | | ✓ | | | | | | | |
| <i>Ochrolechia frigida</i> | | | | ✓ | | | | | | | |
| <i>Ochrolechia pseudopallescens</i> (?) | | | | ✓ | | | | | | | |
| <i>Parmelia omphalodes</i> | | | | ✓ | | | ✓ | | | | |
| <i>Parmelia saxatilis</i> | | | | ✓ | | | | ✓ | | | |
| <i>Parmelia squarrosa</i> | | | ✓ | | | | | ✓ | ✓ | | |
| <i>Parmelia sulcata</i> | ✓ | | | | | | | | | | |
| <i>Parmeliopsis capitata</i> | | | ✓ | | | | | | | | |
| <i>Parmeliopsis hyperopta</i> | | | | ✓ | | | | | | | |
| <i>Peltigera aphthosa</i> | ✓ | | | ✓ | | | | | | | |
| <i>Peltigera canina</i> | ✓ | ✓ | | | | | | | ✓ | | |
| <i>Peltigera degenii</i> | ✓ | | | ✓ | | | ✓ | | | | |
| <i>Peltigera didactyla</i> | ✓ | | | | | | | | | | |
| <i>Peltigera elizabethae</i> | | | | | | ✓ | | | ✓ | | |
| <i>Peltigera membranaceae</i> | ✓ | | | | | ✓ | | | ✓ | | |
| <i>Peltigera praetextata</i> | | | | | | | | | ✓ | | |

| Species | Sir Richard Squires | Stanleyville | Western Brook Pond | Big Level | #2 Lomond River | Burnt Hill | Trout River Trail | Green Garden | Lobster Cove Head | Stuckless Pond | Kildevil Campground |
|----------------------------------|---------------------|--------------|--------------------|-----------|-----------------|------------|-------------------|--------------|-------------------|----------------|---------------------|
| <i>Peltigera scabrosa</i> | | | | ✓ | | | | | | | |
| <i>Pertusaria dactylina</i> | | | | ✓ | | | | | | | |
| <i>Pertusaria macounii</i> | | | | | | | | | | | ✓ |
| <i>Pertusaria panyrga</i> | | | | ✓ | | | | | | | |
| <i>Placynthiella uliginosa</i> | | | | ✓ | | | | | | | |
| <i>Platismatia glauca</i> | ✓ | ✓ | ✓ | ✓ | ✓ | | ✓ | ✓ | ✓ | | |
| <i>Platismatia norvegica</i> | | | | | ✓ | | | | | | |
| <i>Porpidia crustulata</i> | | | | ✓ | | | | | | | |
| <i>Propidia tuberculosa</i> | | | | ✓ | | | | | | | |
| <i>Protopannaria pezizoides</i> | | | | | | | | | ✓ | | |
| <i>Ramalina dilacerata</i> | ✓ | | | | | | | | | | |
| <i>Ramboldia cinnabarina</i> (?) | | ✓ | | | | | | | | | |
| <i>Rhizocarpon geographicum</i> | ✓ | | | ✓ | | | | | | | |
| <i>Ropalospora chlorantha</i> | | | | | | | | | ✓ | | |
| <i>Sphaerophorus fragilis</i> | | | | ✓ | | | | | | | |
| <i>Sphaerophorus globosus</i> | | | | ✓ | | | | | | | |
| <i>Stereocaulon vesuvianum</i> | | | | ✓ | | | | | | | |
| <i>Tuckermannopsis americana</i> | ✓ | | ✓ | | | | | ✓ | | | |
| <i>Tuckermannopsis orbata</i> | | | | | | | | | | | |
| <i>Tuckermannopsis sepincola</i> | | | ✓ | | | | | | | | |
| <i>Umbilicaria cylindrica</i> | | | | ✓ | | | | | | | |
| <i>Umbilicaria hyperborea</i> | | | | ✓ | | | | | | | |
| <i>Umbilicaria polyphylla</i> | | | | ✓ | | | | | | | |
| <i>Umbilicaria proboscidea</i> | | | | ✓ | | | | | | | |
| <i>Usnea dasopoga</i> | ✓ | | | | | | | | | | |
| <i>Usnea longissima</i> | | | ✓ | | | | | | | | |
| <i>Variolaria amara</i> | ✓ | | ✓ | | | | ✓ | | ✓ | | |
| <i>Vulpicida pinastri</i> | | | ✓ | ✓ | | | | ✓ | | | |
| <i>Xanthoparmelia conspersa</i> | | | | | | ✓ | | | | | |
| <i>Xylographa parallela</i> | | ✓ | | | | | | | | | |
| <i>Xylographa</i> sp. | | ✓ | | | | | | | | | |

Note: species in bold may be considered new to the province.

SURVEY OF THE LICHEN-FORMING ASCOMYCETES DURING THE 2015 NL FORAY

Chris Deduke and Michele Piercey-Normore

Lichen collections were made by the authors with the help of A. Arsenault, C. Hanel, and other participants of the foray. This year collections were made at eleven locations throughout Gros Morne National Park. Six of the eleven locations were analysed more closely that had more than 20 species per location. The six locations included **Sir Richard Squires Memorial Provincial Park**, a mixed forest habitat consisting of deciduous (birch) and coniferous trees (spruce, larch and fir). The trail also contains some open bog and moist riparian forest along the Humber River. **Stanleyville trail** was a coniferous forest consisting of mainly balsam fir and decaying wood. **Western Brook Pond trail** had both coniferous forest and large areas of open *Sphagnum* bog. The **Big Level** was a remote arctic/alpine habitat interspersed with balsam fir tuckamore. The **Green Garden trail** was a tableland that ascended into a mixed-wood forest of birch, maple, larch and balsam fir. **Lobster Cove Head trail** was a coastal coniferous forest of fir and spruce with tuckamore.

Species Highlights

There were 124 species or subspecies of lichens reported this year, with 242 specimens collected. Thirty-seven of the species belong to the genus *Cladonia*, five Pin lichens (*Calicium abietinum*, *Calicium* species, *Chaenotheca brunneola*, *C. furfuracea* and *Mycocalicium subtile* (?)), and 10 species of cyanolichens (8 *Peltigera* species, *Protopannaria pezizoides* and *Stereocaulon vesuvianum*). The most abundant species were *Platismatia glauca*, reported in 8 locations, and then *Cladonia pleurota*, *Hypogymnia physodes*, and *H. tubulosa*, which were reported in 6 locations each. Four species were not found on any previous list for the province and may be new to the province, *Calicium abietinum*, *Cetraria ericetorum*, *Hypogymnia krogiae*, and *Ochrolechia pseudopallescens* (?).



Photo: Maria Voik

Lobster Head Cove

Habitat similarity among locations visited

An analysis of habitat similarity (Table 1) between the six locations with over 20 species present shows that Sir Richard Squires Park and Western Brook Pond trail were the most similar with 36.8% (Table 1). Both habitats shared coniferous forest and open bogs. Western Brook Pond trail and Lobster Cove Head trail were the next most similar sites with 30.4% (Table 1). Stanleyville trail shared the most species (28.8%, Table 1) with Sir Richard Squires Park. Both trails had forests populated with birch and spruce trees. Trails in which lichens were not collected are not included in Table 1.

Table 1: Species diversity is shown in this table by the Jaccard's Index of similarity, which is the percent similarity between two locations based on the species composition. The Jaccard's Index of similarity assumes similarity between locations or habitats using the concept that species are adapted to specific habitat conditions. If the composition of species are similar between habitats, then the habitats must also be similar.

| Locations for comparison | Percent similarity |
|---|--------------------|
| Sir Richard Squires X Stanleyville | 28.8% |
| Sir Richard Squires X Western Brook | 36.8% |
| Sir Richard Squires X Big Level | 15.9% |
| Sir Richard Squires X Green Garden | 15.4% |
| Sir Richard Squires X Lobster Cove Head | 27.7% |
| Stanleyville X Western Brook | 21.8% |
| Stanleyville X Big Level | 7.1% |
| Stanleyville X Green Garden | 8.9% |
| Stanleyville X Lobster Cove Head | 19.5% |
| Western Brook X Big Level | 18.6% |
| Western Brook X Green Garden | 22.4% |
| Western Brook X Lobster Cove Head | 30.4% |
| Big Level X Green Garden | 20.0% |
| Big Level X Lobster Cove Head | 5.0% |
| Green Garden X Lobster Cove Head | 10.5% |

The Big Level was the most unique habitat visited at the foray with the lowest average species similarity values between locations (8.4%). The Big Level was most similar to Green Gardens with only 20.0% similarity (Table 1). The two locations shared 14 species, 10 of which were from the genus *Cladonia*. The open barrens from the Tablelands and the arctic/alpine environment offered similar habitat conditions for the ground dwelling lichens. The Big Level is different from Lobster Cove Head at only 5.0% similarity (Table 1). The two locations are different in elevation, forest cover and exposure. The Big Level is an elevated open environment while Lobster Cove Head is a coastal forested habitat.

Green Gardens Trail was the second most distinct habitat, with the second lowest similarity values (11.5%). It shared the highest species similarity with Western Brook Pond at 22.4% (Table 1). Seven of the 11 shared species belonged to the genus *Cladonia* including *C. borealis*, *C. cristatella*, *C. pleurota* and various Reindeer lichens. In general, Western Brook Pond and Sir Richard Squires Park had the highest species similarity values between locations (26.0% and 24.9% respectively), indicating that the forested and bog habitats provided sufficient conditions for a wide variety of different lichen species.

Species Diversity

The Big Level had the highest species diversity with 63 species (Figure 1). The majority of these species were found growing among mosses on the ground. These included members of the *Cladonia*, *Cetraria*, *Ochrolechia* and *Peltigera* genera. Exposed rocks provided additional substrates for lichen growth including foliose members of *Arctoparmelia*, *Melanelia* and *Umbilicaria*, along with crustose *Pertusaria* and *Porpidia* species. Sir Richard Squires Park and Western Brook Pond trail were the next most diverse locations with 39 species each (Figure 1). The *Sphagnum* bog habitat contained many species of *Cladonia*, while the forests were suitable for *Bryoria trichodes*, various *Hypogymnia* species, *Lobaria pulmonaria* and *Platismatia glauca*. Western Brook Pond trail also contained *L. scrobiculata* and *L. quercizans*. Stanleyville trail had a diversity of 28 species, aided in part by the amount of dead and decaying wood in the forest. This provided suitable habitat for crustose species like *Graphis scripta* and *Xylographa parallela*. Green Gardens and Lobster Cove Head each had a diversity of 21 species.



Photo: Andrus Voitk

Nephroma arcticum

Photo: Maria Voik



Graphis scripta




Figure 1: The number of lichen species (y-axis) collected at each location visited (x-axis) during the NL Foray 2015 is shown in this bar graph. The four locations with more than 20 species were compared using Jaccard's Index of similarity.

In summary, this year's foray recorded 124 species of lichens. The Big Level was an exciting new location visited this year, and it provided the greatest lichen diversity with 63 species. This habitat was the most unique of all locations producing the lowest average species similarity scores, comprised primarily of an arctic/alpine environment. The majority of lichens found here were ground dwelling *Cladonia* species and rock dwelling foliose and crustose lichens. The Green Garden trail was the closest in species similarity to the Big Level and this was due to the exposed Tableland barrens at the beginning of the trail. Lobster Head Cove, another new location this year, showed some interesting finds such as *Ropalospora chlorantha* and *Protopannaria pezizoides*.



FORAY NEWFOUNDLAND AND LABRADOR



Photo: Roger Smith

GROS MORNE 2015

OUR PARTNER ORGANIZATIONS

PEOPLE OF NEWFOUNDLAND AND LABRADOR:

DEPARTMENT OF ENVIRONMENT AND CONSERVATION
PARKS AND NATURAL AREAS DIVISION
WILDLIFE DIVISION
FORESTRY AND AGRIFOODS AGENCY
CENTER FOR FOREST SCIENCE AND INNOVATION



PEOPLE OF CANADA, THROUGH

PARKS CANADA

TERRA NOVA NATIONAL PARK
GROS MORNE NATIONAL PARK



THE GROS MORNE CO-OPERATING ASSOCIATION



MEMORIAL UNIVERSITY OF NEWFOUNDLAND

ST. JOHN'S CAMPUS
GRENFELL CAMPUS



TUCKAMORE LODGE



QUIDI VIDI BREWING COMPANY





FORAY

NEWFOUNDLAND AND LABRADOR

2016 2016 2016
2016 2016 2016
2016 2016 2016
2016 2016 2016
2016 2016 2016
2016 2016 2016
2016 2016 2016

The second decade

GOOSE BAY, LABRADOR

Come, visit the Big Land!

September 9-11, 2016

GUEST FACULTY

Renée Lebeuf
Jean Lodge
Michele Piercey-Normore
Roger Smith
Greg Thorn

Get to know our MUSHROOMS & LICHENS!

See our website April/May, 2016, for
Registration Forms & Information:

[<www.nlmushrooms.ca>](http://www.nlmushrooms.ca)

Photo: Muskrat Falls, Labrador, Mavis Penney